

ON EXTREMAL HYPERGRAPHS FOR HAMILTONIAN CYCLES

ROMAN GLEBOV, YURY PERSON, AND WILMA WEPS

ABSTRACT. We study sufficient conditions for Hamiltonian cycles in hypergraphs, and obtain both Turán- and Dirac-type results. While the Turán-type result gives an exact threshold for the appearance of a Hamiltonian cycle in a hypergraph depending only on the extremal number of a certain path, the Dirac-type result yields a sufficient condition relying solely on the minimum vertex degree.

1. INTRODUCTION AND RESULTS

1.1. Turán-type Results. For a fixed graph G we say that the *extremal number* $\text{ex}(n, G)$ of G is the largest integer m such that there exists a graph on n vertices with m edges that does not contain a subgraph isomorphic to G . The corresponding graphs are called *extremal graphs*. Naturally, one can extend this definition to a forbidden spanning structure, e.g. a Hamiltonian cycle (for definition see e.g. [4]). In [15] Ore proved that a non-Hamiltonian graph on n vertices has at most $\binom{n-1}{2} + 1$ edges, and further, that the unique extremal example is given by an $(n-1)$ -clique and a vertex of degree one that is adjacent to one vertex of the clique.

A k -uniform hypergraph H , or k -graph for short, is a pair (V, E) with a vertex set $V = V(H)$ and an edge set $E = E(H) \subseteq \binom{V}{k}$.

There are several definitions of Hamiltonian cycles in hypergraphs, e.g. Berge Hamiltonian cycles [3]. This paper yet follows the definition of Hamiltonian cycles established by Katona and Kierstead [10] as it has become more and more popular in research.

An l -tight Hamiltonian cycle in H , $0 \leq l \leq k-1$, $(k-l)|n$, is a spanning sub- k -graph whose vertices can be cyclically ordered in such a way that the edges are segments of that ordering and every two consecutive edges intersect in exactly l vertices. We denote an l -tight Hamiltonian cycle in a k -graph H on n vertices by $C_n^{(k,l)}$, and call it *tight* if it is $(k-1)$ -tight.

Working on her thesis [25] in coding theory, Woitas raised the question whether removing $\binom{n-2}{2} - 1$ edges from a 3-uniform hypergraph on n vertices leaves a hypergraph containing a 1-tight Hamiltonian cycle. A generalization of this problem is to estimate the extremal number of Hamiltonian cycles in k -graphs.

Katona and Kierstead were the first to study sufficient conditions for the appearance of a $C_n^{(k,k-1)}$ in k -graphs. In [10] they showed that for all integers k and

Date: October 24, 2010.

The first author was supported by DFG within the research training group "Methods for Discrete Structures".

The second author was supported by GIF grant no. I-889-182.6/2005.

n with $2 \leq k$ and $2k - 1 \leq n$,

$$\text{ex}(n, C_n^{(k,k-1)}) \geq \binom{n-1}{k} + \binom{n-2}{k-2}.$$

In the same paper Katona and Kierstead proved, that this bound is not tight for $k = 3$ by showing that for all integers n and q with $q \geq 2$ and $n = 3q + 1$,

$$\text{ex}(n, C_n^{(3,2)}) \geq \binom{n-1}{3} + n - 1.$$

In [24] Tuza gave a construction for general k and tight Hamiltonian cycles, improving the lower bound to

$$\text{ex}(n, C_n^{(k,k-1)}) \geq \binom{n-1}{k} + \binom{n-1}{k-2},$$

if a Steiner system $S(k-2, 2k-3, n-1)$ exists. Also for all k, n and p such that a partial Steiner system $PS(k-2, 2k-3, n-1)$ of order $n-1$ with $p \binom{n-1}{k-2} / \binom{2k-3}{k-2}$ blocks exists, Tuza proved the bound

$$\text{ex}(n, C_n^{(k,k-1)}) \geq \binom{n-1}{k} + p \binom{n-1}{k-2}.$$

An intuitive approach to forbid Hamiltonian cycles in hypergraphs is to prohibit certain structures in the *link* of one fixed vertex. For a vertex $v \in V$, we define the *link of v in H* to be the $(k-1)$ -graph $H(v) = (V \setminus \{v\}, E_v)$ with $\{x_1, \dots, x_{k-1}\} \in E_v$ iff $\{v, x_1, \dots, x_{k-1}\} \in E(H)$.

The structure of interest in this case is the generalization of a path for hypergraphs.

An l -tight k -uniform t -path, denoted by $P_t^{(k,l)}$, is a k -graph on t vertices, $(k-l)|(t-l)$, such that there exists an ordering of the vertices, say (x_1, \dots, x_t) , in such a way that the edges are segments of that ordering and every two consecutive edges intersect in exactly l vertices. Observe that a $P_t^{(k,l)}$ has $\frac{t-l}{k-l}$ edges. A k -uniform $(k-1)$ -tight path is called *tight*, and whenever we consider a path we assume it to be tight unless stated otherwise.

For arbitrary k and l we give the exact extremal number and the extremal graphs of l -tight Hamiltonian cycles in this paper. The extremal number and the extremal graphs rely on the extremal number of $P(k, l) := P_{\lfloor \frac{k}{k-l} \rfloor}^{(k-1, l-1)}(k-l)+l-1$, and its extremal graphs, respectively.

Theorem 1. *For any $k \geq 2, l \in \{0, \dots, k-1\}$ there exists an n_0 such that for any $n \geq n_0$ and $(k-l)|n$,*

$$\text{ex}\left(n, C_n^{(k,l)}\right) = \binom{n-1}{k} + \text{ex}(n-1, P(k, l))$$

holds. Furthermore, any extremal graph on n vertices contains an $(n-1)$ -clique and a vertex whose link is $P(k, l)$ -free.

Actually, in this paper we prove a stronger statement, namely that with one more edge we find a Hamiltonian cycle that is l -tight in the neighborhood of one vertex and is $(k-1)$ -tight on the rest.

Using the result by Györi, Katona and Lemons [7] stating that

$$(1 + o(1)) \binom{n-1}{k-2} \leq \text{ex}(n-1, P_{2k-2}^{(k-1, k-2)}) \leq (k-1) \binom{n-1}{k-2},$$

we obtain lower and upper bounds for $l = k - 1$:

$$\binom{n-1}{k} + (1 + o(1)) \binom{n-1}{k-2} \leq \text{ex}(n, C_n^{(k, k-1)}) \leq \binom{n-1}{k} + (k-1) \binom{n-1}{k-2}.$$

Note that the upper bound also holds for $l \neq k - 1$.

Furthermore, for $k = 3$ and $l = 2$ Theorem 1 states that there exists an n_0 such that for any $n \geq n_0$,

$$\begin{aligned} \text{ex}(n, C_n^{(3,2)}) &= \binom{n-1}{3} + \text{ex}(n-1, P_4^{(2,1)}) \\ &= \begin{cases} \binom{n-1}{3} + n - 1, & 3 \mid n - 1 \\ \binom{n-1}{3} + n - 2, & \text{otherwise.} \end{cases} \end{aligned}$$

In our proof we make use of the *absorbing technique* that was originally developed by Rödl, Ruciński and Szemerédi.

1.2. Dirac-type Results. The problem of finding Hamiltonian cycles and perfect matchings in graphs has been studied very intensively. There are plenty beautiful conditions guaranteeing the existence of such cycles, e.g. Dirac's condition [6].

Over the last couple of years several Dirac-type results in hypergraphs were shown, and along with them, different definitions of *degree* in a k -graph were introduced. They all can be captured by the following definition. The *degree* of $\{x_1, \dots, x_i\}$, $1 \leq i \leq k - 1$, in a k -graph H is the number of edges the set is contained in and is denoted by $\deg(x_1, \dots, x_i)$. Let

$$\delta_d(H) := \min\{\deg(x_1, \dots, x_d) \mid \{x_1, \dots, x_d\} \subset V(H)\}$$

for $0 \leq d \leq k - 1$. If the graph is clear from the context, we omit H and write for short δ_d . Note that $\delta_0 = e(H) := |E(H)|$ and δ_1 is the minimum vertex degree in H .

Following the definitions of Rödl and Ruciński in [17], denote for every d, k, l and n with $0 \leq d \leq k - 1$ and $(k - l) \mid n$ the number $h_d^l(k, n)$ to be the smallest integer h such that every n -vertex k -graph H satisfying $\delta_d(H) \geq h$ contains a l -tight Hamiltonian cycle. Observe that $h_0^l(k, n) = \text{ex}(n, C_n^{(k,l)}) + 1$.

In [10] Katona and Kierstead showed that $h_{k-1}^{k-1}(k, n) \geq \lfloor \frac{n-k+3}{2} \rfloor$ by giving an extremal construction. Their implicit conjecture that this bound is tight was confirmed for $k = 3$ by Rödl, Ruciński and Szemerédi in [19] asymptotically and in [23] exactly. For $k \geq 4$ the same authors showed in [21] that $h_{k-1}^{k-1}(k, n) \sim \frac{1}{2}n$. Generalizing the results to other tightnesses, Markström and Ruciński proved in [14] that $h_{k-1}^l(k, n) \sim \frac{1}{2}n$ if $(k - l) \mid k, n$. In [11] Kühn, Mycroft and Osthus proved that

$$h_{k-1}^l(k, n) \sim \frac{n}{\left\lceil \frac{k}{k-l} \right\rceil (k-l)}$$

if $k - l$ does not divide k and $(k - l) \mid n$, proving a conjecture by Hàn and Schacht [9]. For further information, an excellent survey of the recent results can be found in [17].

Rödl and Ruciński conjectured in [17] that for all $1 \leq d \leq k-1$, $k|n$,

$$h_d^{k-1}(k, n) \sim h_d^0(k, n).$$

Further notice that 0-tight Hamiltonian cycles $C_n^{(k,0)}$ are perfect matchings. Perfect matching may be considered the “simplest” spanning structure and there are several results about $h_{k-1}^0(k, n)$, see [20, 12, 22, 18, 1, 16, 8, 13].

Noting the fact that there are virtually no results on $h_d^l(k, n)$ for $d \leq k-2$, Rödl and Ruciński remarked in [17] that it does not even seem completely trivial to show $h_1^2(3, n) \leq c \binom{n-1}{2}$ for some constant $c < 1$. Further, they gave the following bounds

$$\left(\frac{5}{9} + o(1)\right) \binom{n-1}{2} \leq h_1^2(3, n) \leq \left(\frac{11}{12} + o(1)\right) \binom{n-1}{2}.$$

We show the following upper bound on $h_1^{k-1}(k, n)$.

Theorem 2. *For any $k \in \mathbb{N}$ there exists an n_0 such that every k -graph H on $n \geq n_0$ vertices with $\delta_1 \geq \left(1 - \frac{1}{22(1280k^3)^{k-1}}\right) \binom{n-1}{k-1}$ contains a tight Hamiltonian cycle.*

Note that Theorem 2 implies

$$h_d^l(k, n) \leq \left(1 - \frac{1}{22(1280k^3)^{k-1}}\right) \binom{n-d}{k-d}$$

for all $l \in \{0, \dots, k-1\}$ and all $1 \leq d \leq k-1$.

2. PROOFS

2.1. Outline of the Proofs. In the following we give a brief overview over the structure of the proofs of Theorems 1 and 2. For this section we define

$$\epsilon = \frac{1}{22(1280k^3)^{k-1}}$$

and

$$\varrho = (22\epsilon)^{\frac{1}{k-1}}.$$

Suppose $H = (V, E)$ is a k -graph on n vertices with $\delta_1 \geq (1 - \epsilon) \binom{n-1}{k-1}$ and n sufficiently large. With an *end* of a path $P_t^{(k,l)}$ we mean its first $k-1$ vertices, (x_1, \dots, x_{k-1}) , or its last $k-1$ vertices in reverse order, (x_t, \dots, x_{t-k+2}) , considering the ordered vertices. For an i -tuple (x_1, \dots, x_i) in H we write \mathbf{x}_i , $1 \leq i \leq n$. We call \mathbf{x}_{k-1} *good* if for all $i \in \{1, \dots, k-1\}$ it holds that

$$\deg(x_1, \dots, x_i) \geq (1 - \varrho^{k-i}) \binom{n-i}{k-i}. \quad (1)$$

A path is called *good* if both of its ends are good.

Outline of the proofs:

1. In Claim 8 we prove the existence of one l -tight good path or several vertex-disjoint good tight paths containing the vertices of small degree. (Note that we do not need this step in the proof of Theorem 2.)

2. We say that a tuple \mathbf{x}_{2k-2} *absorbs* a vertex $v \in V$ if both \mathbf{x}_{2k-2} and $(x_1, \dots, x_{k-1}, v, x_k, \dots, x_{2k-2})$ induce good paths in H , meaning that the corresponding ordering of the paths is \mathbf{x}_{2k-2} or $(x_1, \dots, x_{k-1}, v, x_k, \dots, x_{2k-2})$, respectively, and the ends are good. Lemma 4 ensures a set \mathcal{A} , providing that any vertex remaining can be absorbed by many tuples of \mathcal{A} . We call an element of \mathcal{A} *absorber*.
3. For $\mathbf{x}_i, \mathbf{y}_j \in V^{k-1}$ we define

$$\mathbf{x}_i \times \mathbf{y}_j := (x_1, \dots, x_i, y_1, \dots, y_j).$$

Let \mathbf{x}_{k-1} and \mathbf{y}_{k-1} be good. We say that a tuple \mathbf{z}_{k-1} *connects* \mathbf{x}_{k-1} with \mathbf{y}_{k-1} if $(x_{k-1}, \dots, x_1) \times \mathbf{z}_{k-1} \times \mathbf{y}_{k-1}$ induces a path in H with respect to the order. Notice that the connecting-operation is not symmetric. Lemma 5 guarantees a set \mathcal{C} such that any pair of $(k-1)$ -tuples in H can be connected by many elements of \mathcal{C} . We call the elements of \mathcal{C} *connectors*.

4. We modify \mathcal{A} and \mathcal{C} such that \mathcal{A} , \mathcal{C} and the element(s) of Step 1 are pairwise vertex-disjoint.
5. In Lemma 6 we create a good tight path that contains all elements of the modified \mathcal{A} , respecting their ordering.
6. Using Lemma 7, we extend the path from Step 5 until it covers almost all of the remaining vertices that neither participate in (l -tight or tight) good paths of Step 1 nor in the modified \mathcal{C} .
7. Using connectors, we create a cycle containing the (l -tight or tight) good paths from Step 1 and the good path from Step 6.
8. In the final step all remaining vertices are absorbed by the absorbers of the cycle.

2.2. Auxiliary Lemmas. In this part we derive the main tools used to prove Theorems 1 and 2. For this subsection let $H = (V, E)$ be a k -graph on n vertices with

$$\delta_1 \geq (1 - \epsilon) \binom{n-1}{k-1}, \quad (2)$$

and recall $\epsilon = \frac{1}{22(1280k^3)^{k-1}}$.

Lemma 3. *Let \mathbf{x}_{2k-2} be chosen u.a.r. from V^{2k-2} . The probability that all x_i s are pairwise distinct and both (x_1, \dots, x_{k-1}) and (x_{2k-2}, \dots, x_k) are good is at least $\frac{8}{11}$.*

Proof. Let a be the number of $(k-1)$ -tuples that are not good and b_j the number of j -tuples \mathbf{y}_j with $\deg(y_1, \dots, y_j) < (1 - \varrho^{k-j}) \binom{n-j}{k-j}$. Thus, by the definition of a good tuple, for each tuple \mathbf{y}_k that is not good, there exists a j such that \mathbf{y}_j is one of the b_j tuples with small degree. Furthermore, for every \mathbf{y}_j there are at most $\frac{(n-j)!}{(n-k+1)!}$ different $(k-1)$ -tuples $(y_1, \dots, y_j, z_1, \dots, z_{k-1-j})$ with pairwise distinct $z_j \in V \setminus \{y_1, \dots, y_j\}$. Hence,

$$a \leq \sum_{j=1}^{k-1} \frac{(n-j)!}{(n-k+1)!} b_j.$$

The second time we apply double counting, we recall that H has at most $\epsilon \binom{n}{k}$ non-edges. Each of the b_j j -tuples is by definition in at least $\varrho^{k-j} \binom{n-j}{k-j}$ non-edges,

and from every non-edge one obtains $\binom{k}{j}j!$ different j -tuples. Thus,

$$\varrho^{k-j} \binom{n-j}{k-j} b_j \leq \binom{k}{j} j! \epsilon \binom{n}{k}.$$

Putting the two bounds together, we obtain

$$\begin{aligned} \Pr[\mathbf{x}_{k-1} \text{ is not good}] &= \frac{a}{n^{k-1}} \leq \sum_{j=1}^{k-1} \frac{(n-j)!}{(n-k+1)!} b_j \frac{1}{n^{k-1}} \\ &\leq \sum_{j=1}^{k-1} \frac{(n-j)!}{(n-k+1)!} \frac{\binom{k}{j} j! \epsilon \binom{n}{k}}{\varrho^{k-j} \binom{n-j}{k-j}} \frac{1}{n^{k-1}} \leq \epsilon \sum_{i=1}^{k-1} \frac{1}{\varrho^{k-i}} < \frac{2\epsilon}{\varrho^{k-1}} = \frac{1}{11}. \end{aligned}$$

Let \mathbf{x}_{2k-2} be chosen u.a.r. from V^{2k-2} . Then

$$\begin{aligned} \Pr[(x_1, \dots, x_{k-1}) \text{ and } (x_{2k-2}, \dots, x_k) \text{ are good and the } x_i\text{s are pairwise distinct}] \\ \geq \frac{\binom{n}{2k-2}}{n^{2k-2}} - \frac{2}{11} \geq 1 - \frac{3}{11} = \frac{8}{11}. \end{aligned}$$

□

For a given set \mathcal{X} of tuples or graphs, we write X when considering the corresponding vertex set.

Lemma 4. *For all γ , $0 < \gamma \leq \frac{1}{64k^2}$, there exists a set \mathcal{A} of size at most $2\gamma n$ consisting of disjoint $(2k-2)$ -tuples, each inducing a good path with respect to its order, such that for each vertex $v \in V$ at least $\frac{\gamma n}{4}$ tuples in \mathcal{A} absorb v .*

Proof. By Lemma 3, we know that there are at least $\frac{8}{11}n^{2k-2}$ tuples $\mathbf{x}_{2k-2} \in V^{2k-2}$, such that the x_i s are pairwise distinct and both (x_1, \dots, x_{k-1}) and (x_{2k-2}, \dots, x_k) are good. We denote the set of such tuples by \mathcal{A}' .

Let v be a vertex from V and denote by \mathcal{A}_v the set of tuples \mathbf{x}_{2k-2} from \mathcal{A}' such that, in addition,

- $\{x_j, \dots, x_{j+k-1}\} \in E(H)$, $1 \leq j \leq k-1$, and
- $\{v, x_j, \dots, x_{j+k-2}\} \in E(H)$, $1 \leq j \leq k$.

Therefore, the set \mathcal{A}_v consists of those paths that can absorb the vertex v . From the minimum degree condition on H , see (2), it follows that

$$|\mathcal{A}_v| \geq \frac{8}{11}n^{2k-2} - 2k\epsilon n^{2k-2} \geq \frac{7}{11}n^{2k-2}. \quad (3)$$

Fix γ with $0 < \gamma \leq \frac{1}{64k^2}$. Let \mathcal{A} be the set obtained by choosing each $(2k-2)$ -tuple $\mathbf{x}_{2k-2} \in V^{2k-2}$ from \mathcal{A}' independently with probability $\frac{\gamma}{n^{2k-3}}$.

The expected size of $|\mathcal{A}|$ is at most γn and we apply Chernoff's inequality (e.g. see [2]):

$$\Pr[|\mathcal{A}| - \gamma n > \gamma n] < e^{-\gamma n}. \quad (4)$$

This way, with high probability we obtain at most $2\gamma n$ many $(2k-2)$ -tuples.

Let Y be the random variable taking the value 1 whenever a pair of tuples in \mathcal{A} is not vertex-disjoint and the value 0 else. Thus,

$$\mathbb{E}[Y] \leq (2k-2)^2 n^{4k-5} \frac{\gamma^2}{n^{4k-6}} \leq 4k^2 \gamma^2 n. \quad (5)$$

Applying Markov's inequality, we obtain:

$$\Pr [Y > 8k^2\gamma^2n] < \frac{1}{2}. \quad (6)$$

From (3) we infer by Chernoff's inequality that

$$\Pr \left[|\mathcal{A}_v \cap \mathcal{A}| < \frac{\gamma n}{2} \right] < e^{-\frac{1}{100}\gamma n} \quad (7)$$

since $\mathbb{E}[|\mathcal{A}'_v \cap \mathcal{A}|] \geq \frac{7}{11}\gamma n$.

With (3), (6) and (7), we see that if we delete from \mathcal{A} those tuples that have vertices in common, we obtain with probability at least

$$1/2 - e^{-\gamma n} - ne^{-\frac{1}{100}\gamma n} > 1/4$$

a new set satisfying the conditions of the lemma (note that $\frac{\gamma n}{2} - 8k^2\gamma^2n \geq \frac{\gamma n}{4}$). \square

The following lemma provides us with the essential tool to close the cycle.

Lemma 5. *For all β , $0 < \beta \leq \frac{1}{64k^2}$, there exists a set \mathcal{C} of size at most $2\beta n$ consisting of pairwise disjoint $(k-1)$ -tuples, such that for each pair of good $(k-1)$ -tuples there exist at least $\frac{\beta n}{4}$ elements in \mathcal{C} that connect this pair.*

Proof. For two good vertex-disjoint tuples $\mathbf{x}_{k-1}, \mathbf{y}_{k-1}$ in H , let $\mathcal{C}_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}}$ be the set of all connectors that connect \mathbf{x}_{k-1} with \mathbf{y}_{k-1} and vertex disjoint from $\mathbf{x}_{k-1}, \mathbf{y}_{k-1}$. Recall that the following conditions hold for $z \in \mathcal{C}_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}}$:

- $\{x_{k-i}, \dots, x_1, z_1, \dots, z_i\} \in E(H)$, for $i \in \{1, \dots, k-1\}$, and
- $\{z_{i-k+1}, \dots, z_{k-1}, y_1, \dots, y_{i-k+1}\} \in E(H)$, for $i \in \{k, \dots, 2k-2\}$.

From the condition (1) from the definition of good tuples and from (2), the minimum degree of H , we infer

$$|\mathcal{C}_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}}| \geq \left(1 - 2 \sum_{i=1}^{k-1} \rho^i\right) n^{k-1} - o(n^{k-1}) \geq (1 - 4\rho k)n^{k-1}.$$

Now, take β as asserted by the lemma and let $\mathcal{C}' := \cup_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}} \mathcal{C}_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}}$, where the union is over all vertex-disjoint good $(k-1)$ -tuples. Define \mathcal{C} to be the set obtained by choosing each $\mathbf{z}_{k-1} \in \mathcal{C}'$ independently with probability $\frac{\beta}{n^{k-2}}$.

Similarly to (4), with high probability $|\mathcal{C}| \leq 2\beta n$. With probability at least $\frac{1}{2}$ at most $4k^2\beta^2n \leq \beta n/4$ of the $(k-1)$ -tuples have to be removed from \mathcal{C} to obtain a set of vertex-disjoint tuples, analogously to (6).

Analogously to (7), for two good vertex-disjoint tuples $\mathbf{x}_{k-1}, \mathbf{y}_{k-1}$ in H ,

$$\Pr \left[|\mathcal{C}_{\mathbf{x}_{k-1}, \mathbf{y}_{k-1}} \cap \mathcal{C}| < \frac{\beta n}{2} \right] < e^{-\frac{1}{16}\gamma n}.$$

Therefore, we deduce with positive probability that removing from \mathcal{C} all tuples that are not vertex-disjoint, we are left with a set that satisfies the conditions in the lemma. \square

The next lemma helps us to connect linear amount of small paths into a single path avoiding a small forbidden vertex subset.

Lemma 6. *For any set \mathcal{X} of vertex-disjoint $(2k-2)$ -tuples that each induce a good path in H , $|\mathcal{X}| \leq \frac{1}{4k^2}n$, and any forbidden set $F \subset V$ of size at most $\frac{1}{8k}n$, there exists a path P containing all tuples of \mathcal{X} , respecting their individual ordering, such that $(V(P) \setminus X) \cap F = \emptyset$.*

Proof. For arbitrary $\mathbf{x}_{2k-2}, \mathbf{y}_{2k-2} \in \mathcal{X}$, we choose a $\mathbf{z}_{k-1} \in V^{k-1}$ uniformly at random and define the events

$$E^1 = \{\mathbf{x}_{2k-2} \times \mathbf{z}_{k-1} \text{ induces a path, respecting the ordering}\}$$

and

$$E^2 = \{\mathbf{z}_{k-1} \times \mathbf{y}_{2k-2} \text{ induces a path, respecting the ordering}\}.$$

With E_i^2 being the event that $\{z_i, \dots, z_{k-1}, y_1, \dots, y_i\} \in E$ we obtain that

$$\Pr[E_i^2] \geq 1 - \varrho^{k-i} - o(1),$$

since \mathbf{x}_{2k-2} and \mathbf{y}_{2k-2} induce good paths and the probability that at least two of the k vertices coincide is $o(1)$. Therefore,

$$\Pr[E^2] \geq 1 - \sum_{i=1}^{k-1} (1 - \Pr[E_i^2]) \geq 1 - \sum_{i=1}^{k-1} (\varrho^{k-i} + o(1)) \geq 1 - 2\varrho.$$

The same holds for E^1 . Hence, by the union bound

$$\Pr[E^1 \cap E^2] \geq 1 - 4\varrho.$$

We choose an arbitrary ordering of \mathcal{X} . Iteratively, we consider two consecutive elements $\mathbf{x}_{2k-2}, \mathbf{y}_{2k-2}$ of \mathcal{X} . The probability that a u.a.r. chosen $\mathbf{z}_{k-1} \in V^{k-1}$ connects \mathbf{x}_{2k-2} with \mathbf{y}_{2k-2} (meaning that both E^1 and E^2 hold) is at least $1 - 4\varrho$ and the probability that it is not vertex-disjoint to an already chosen element (connecting previous pairs of elements of \mathcal{X}), to X or to F is at most

$$\left(\frac{k}{4k^2} + \frac{2k}{4k^2} + \frac{1}{8k}\right)k = \frac{7}{8} < 1 - 4\varrho$$

by the union bound.

Thus, we choose a $\mathbf{z}_{k-1} \in V^{k-1}$ satisfying the conditions in the lemma, and iterate. \square

The next lemma helps us find an almost spanning path in the hypergraph H .

Lemma 7. *For every good path P , there exists a set $F \subset V$ of size at most $k\varrho n$ such that P can be extended to a longer good path with any vertex from $V \setminus (F \cup V(P))$.*

Proof. Consider the longest good path P' that contains P and suppose that $|V(P')| < n - k\varrho n$. Then choose one end \mathbf{x}_{k-1} of P' . Note that from (1), i.e., from $\deg(x_1, \dots, x_i) \geq (1 - \varrho^{k-i}) \binom{n-i}{k-i}$ for every i , it follows that, for every i , the number of vertices $v \in V(H)$ such that

$$\deg(v, x_1, \dots, x_i) \geq (1 - \varrho^{k-i-1}) \binom{n-i-1}{k-i-1} \tag{8}$$

is at least ϱn , implying that $|P'| \geq n - k\varrho n$.

Indeed, suppose for contradiction that there exists an i , such that the number of vs satisfying (8) is less than $n - \varrho n$. Then,

$$\begin{aligned} \deg(x_1, \dots, x_i) &= \sum_{v \in V \setminus \{x_1, \dots, x_i\}} \frac{1}{k-i} \deg(v, x_1, \dots, x_i) < \\ &\frac{(n - \varrho n)}{k-i} \binom{n-i-1}{k-i-1} + \frac{\varrho n - i}{k-i} (1 - \varrho^{k-i-1}) \binom{n-i-1}{k-i-1} \leq (1 - \varrho^{k-i}) \binom{n-i}{k-i}, \end{aligned}$$

contradicting (1). \square

3. PROOFS OF THEOREMS 1 AND 2

Proof. Suppose $H = (V, E)$ is a k -graph with at least

$$\binom{n-1}{k} + \text{ex}(n-1, P(k, l))$$

edges and no vertex with a $P(k, l)$ -free link. Then the vertex set can be partitioned into two sets $V = V' \cup V''$ with $|V'| = n'$ and $V'' = \{v_1, \dots, v_t\}$ such that

$$\delta_1(H') \geq (1 - \epsilon) \binom{n'}{k-1} \quad (9)$$

with $H' = H[V']$. To obtain V' , we iteratively delete vertices v_1, \dots, v_t of minimum degree from H till the δ_1 -condition (9) holds. Counting the non-edges one observes that $t \leq \frac{2}{\epsilon}$.

The following claim provides an embedding of the vertices of V'' .

Claim 8. *There exists a set \mathcal{S} of t paths of type $P_{2k-2}^{(k, k-1)}$, if $t \geq 2$, or of type $P(k, l)$, if $t = 1$, such that v_i is in each edge of the i^{th} element of \mathcal{S} , $1 \leq i \leq t$, for some ordering of \mathcal{S} .*

We apply Lemma 4 for $\gamma = \frac{1}{64k^2}$ and Lemma 5 for $\beta = \frac{1}{1280k^3}$ to H' . As we want disjoint sets \mathcal{A} , \mathcal{C} and \mathcal{S} , we delete all elements from \mathcal{A} that are not vertex-disjoint to an element from \mathcal{C} or \mathcal{S} . Thus, we delete at most $2k\beta n' + \frac{4k}{\epsilon} \leq \frac{\gamma n'}{20}$ absorbers overall, and for every vertex $v \in V'$ there are at least $\frac{\gamma n'}{5}$ elements in the new set \mathcal{A} absorbing v . Similarly, we make \mathcal{C} disjoint from \mathcal{S} still keeping at least $\frac{\beta n'}{5}$ connectors in \mathcal{C} for each pair of good $(k-1)$ -tuples.

Applying Lemma 6 on the new set \mathcal{A} , we obtain a good tight path in H' containing all elements of \mathcal{A} and no vertex from $\mathcal{C} \cup \mathcal{S}$. We extend this path to one good path with Lemma 7 such that it covers all but $k\alpha n'$ vertices from $V' \setminus (\mathcal{C} \cup \mathcal{S})$ and does not contain any vertex from $\mathcal{C} \cup \mathcal{S}$.

As a next step, we use connectors from \mathcal{C} to connect the elements of \mathcal{S} and the extended path to one cycle. This cycle absorbs the remaining vertices including the unused connectors, since $2\beta n' + k\alpha n' < \frac{\gamma n'}{5}$. If \mathcal{S} contains only one element, we obtain a Hamiltonian cycle that is tight except for the l -tight path from \mathcal{S} . Otherwise, we obtain a tight Hamiltonian cycle. Hence, there exists an l -tight Hamiltonian cycle. \square

Note that we actually prove the bound for Hamiltonian cycles that are tight except in the link of at most one vertex.

Now deliver the missing proof of the above claim.

Proof of the Claim 8. We consider two cases.

Case 1 ($\deg(v_1) < \frac{\epsilon}{2} \binom{n-1}{k-1}$). In this case, the number of missing edges yields a sufficient minimum degree in $H - v_1$, hence $t = 1$.

Let a be the number of hyperedges $\{x_1, \dots, x_{k-1}\}$ in $H(v_1)$ that contain a subset $\{x_1, \dots, x_j\}$, $j \in \{1, \dots, k-1\}$, satisfying $\deg(x_1, \dots, x_j) < (1 - \rho^{k-j}) \binom{n-1-j}{k-j}$, i.e., if there is a tuple (x_1, \dots, x_{k-1}) obtained by an ordering of the edge that is not good in H' . We denote the number of such j -sets by b_j and observe that each of

them lies in at most $\binom{n-1-j}{k-1-j}$ edges in $H(v_1)$. Hence, we obtain

$$a \leq \sum_{j=1}^{k-1} \binom{n-1-j}{k-1-j} b_j.$$

We further call those a edges *bad*.

The second time we apply double counting, we set c to be the number of non-edges in H' . By definition, each of the b_j j -sets lies in at least $\varrho^{k-j} \binom{n-1-j}{k-j}$ non-edges of H' . Note that each non-edge of H' has exactly 2^k subsets. Henceforth,

$$\sum_{j=1}^{k-1} \varrho^{k-j} \binom{n-1-j}{k-j} b_j \leq 2^k c.$$

Combining the two bounds, there are at most

$$\begin{aligned} a &\leq \sum_{j=1}^{k-1} \binom{n-1-j}{k-1-j} b_j \leq \sum_{j=1}^{k-1} \varrho^{k-j} \binom{n-1-j}{k-j} b_j \frac{k-j}{n-k} \varrho^{j-k} \\ &\leq \sum_{j=1}^{k-1} \varrho^{k-j} 2^k c \frac{k-j}{n-k} \varrho^{j-k} \leq 2^{k+1} c \frac{k-1}{n-k} \varrho^{1-k} < c \end{aligned}$$

edges in $H(v_1)$, which have an ordering producing a tuple that is not good in H' . Observe that equality can only be obtained with $c = 0$.

For $c = 0$, there exists a $P(k, l)$ in the link of v_1 per definition of H , and this path is good, hence, we are done. For $c > 0$, we obtain $\deg(v_1) > c + \text{ex}(n-1, P(k, l))$. We disregard *bad* hyperedges in $H(v_1)$ and using $a < c$, we still find a $P(k, l)$ in the link of v_1 . The obtained path is good, proving the lemma.

Case 2 ($\deg(v_1) \geq \frac{\epsilon}{2} \binom{n-1}{k-1}$). Here we have for all $1 \leq i \leq t$, $\deg(v_i) \geq \frac{1}{3} \binom{n-1}{k-1}$ holds because the v_i s are chosen greedily with ascending degree. In this case, we actually show that each of the v_i s can be matched to a good tight path such that the assigned paths are pairwise vertex-disjoint. Since the edge density of H' is $1 - o(1)$, we know that there are $o(1) \binom{n'}{k-1}$ tuples that are not good in H' . By the result from [7] mentioned in the introduction it holds that

$$\text{ex}\left(n', P_{2k-2}^{(k-1, k-2)}\right) \leq (k-1) \binom{n'}{k-2} = o(1) \binom{n'}{k-1}.$$

There are at most $O(1) \binom{n-2}{k-2}$ edges including at least two vertices from V'' . We assign iteratively vertex-disjoint good $(2k-1)$ -paths to each of the v_i , $1 \leq i \leq t$, such that v_i is in each of its edges. This is possible, since we disregard at most $o(1) \binom{n'}{k-1}$ many edges in the link of each v_i that contain a tuple that is not good or a vertex contained in a previously assigned path or another vertex from V'' .

□

Proof of Theorem 2. The proof of Theorem 2 follows the same pattern as the proof of Theorem 1 without making use of Claim 8. Therefore, we only give a brief sketch of it.

Suppose H is a k -graph with $\delta_1 \geq (1 - \epsilon) \binom{n-1}{k-1}$. Similarly to the proof of Theorem 1, we apply Lemmas 4 and 5 and obtain via deletion of elements of \mathcal{A} two vertex-disjoint sets A and C such that \mathcal{A} and C have the desired properties. Using

Lemma 6 we find a good path containing all elements of \mathcal{A} such that \mathcal{C} is disjoint to it. We extend this path with Lemma 7 such that it contains all but at most ρkn vertices from $V \setminus \mathcal{C}$ and no vertex from \mathcal{C} . Using a connector, we connect the ends of this path, obtaining a cycle. As $2\beta n + k\rho n < \frac{2n}{5}$ holds, we absorb the remaining vertices and obtain a tight Hamiltonian cycle. \square

4. CONCLUDING REMARKS

The asymptotic edge-density of extremal non-Hamiltonian hypergraphs is $1 - o(1)$ (unlike the density of F -extremal graphs for fixed k -graphs F), since a Hamiltonian cycle is a spanning substructure. In general, we conjecture that an extremal graph of any bounded spanning structure consists of an $(n-1)$ -clique and a further extremal graph.

Conjecture 9. *For any $k \in \mathbb{N}$ there exists an n_0 such that for every k -graph H on $n \geq n_0$ vertices without a spanning subgraph isomorphic to a forbidden hypergraph F of bounded maximum vertex degree,*

$$|e(H)| \leq \binom{n-1}{k} + \text{ex}(n-1, \{F(v) : v \in V\})$$

holds, and the bound is tight.

It is not hard to see that Conjecture 9 holds for forbidden spanning subgraphs F containing a fixed vertex set $V' \subset V$, where $|V'|$ has constant size, such that $F[V \setminus V']$ is a subgraph consisting of a vertex-disjoint union of paths. Note that Conjecture 9 implies the result obtained by the proof of Theorem 1.

A 2-graph is called *pancyclic*, if for any c with $3 \leq c \leq n$ it contains a c -cycle. Similarly to the spanning structure of Hamiltonian l -tight cycles, Katona and Kierstead [10] defined l -tight cycles of any length. This allows us to generalize the concept of pancyclicity by calling a k -graph *l -pancyclic*, if for any c with $3 \leq c \leq n/(k-l)$ it contains an l -tight cycle on c edges. In his famous metaconjecture in [5], Bondy noted for 2-graphs that almost any non-trivial condition on a graph which implies that the graph is Hamiltonian also implies that the graph is pancyclic. (There may be a simple family of exceptional graphs.)

It is not hard to see that both the condition in Theorem 1 and the condition in Theorem 2 imply not only Hamiltonicity but also pancyclicity.

REFERENCES

- [1] R. Aharoni, A. Georgakopoulos, and P. Sprüssel. Perfect matchings in r -partite r -graphs. *Eur. J. Comb.*, 30(1):39–42, 2009. 1.2
- [2] N. Alon and J. H. Spencer. *The probabilistic method. With an appendix on the life and work of Paul Erdős. 3rd ed.*, pages 307–313. Wiley-Interscience Series in Discrete Mathematics and Optimization. Hoboken, NJ: John Wiley & Sons, 2008. 2.2
- [3] J.C. Bermond, A. Germa, M.C. Heydemann, and D. Sotteau. Hypergraphes hamiltoniens. *Problèmes combinatoires et théorie des graphes, Orsay 1976, Colloq. int. CNRS*, 260:39–43, 1978. 1.1
- [4] Bollobás, B. *Modern graph theory*, pages 14–16. Graduate Texts in Mathematics. 184. New York, NY: Springer. xiii, 1998. 1.1
- [5] J.A. Bondy. Pancyclic graphs: recent results. *Infinite and finite Sets, Colloq. Honour Paul Erdős, Keszthely 1973, Colloq. Math. Soc. Janos Bolyai 10*, 181–188, 1975. 4
- [6] G.A. Dirac. Some theorems on abstract graphs. *Proc. Lond. Math. soc., III. Ser.*, 2:69–81, 1952. 1.2

- [7] E. Györi, G. Y. Katona, and N. Lemons. Hypergraph extensions of the Erdős-Gallai theorem. *Discrete Math.*, 36:655–662, 2010. [1.1](#), [2](#)
- [8] H. Hàn, Y. Person, and M. Schacht. On perfect matchings in uniform hypergraphs with large minimum vertex degree. *SIAM J. Discrete Math.*, 23(2):732–748, 2009. [1.2](#)
- [9] H. Hàn and M. Schacht. Dirac-type results for loose Hamilton cycles in uniform hypergraphs. *J. Comb. Theory, Ser. B*, 100(3):332–346, 2010. [1.2](#)
- [10] G.Y. Katona and H.A. Kierstead. Hamiltonian chains in hypergraphs. *J. Graph Theory*, 30(3):205–212, 1999. [1.1](#), [1.2](#), [4](#)
- [11] D. Kühn, R. Mycroft, and D. Osthus. Hamilton ℓ -cycles in uniform hypergraphs. *J. Comb. Theory, Ser. A*, 117(7):910–927, 2010. [1.2](#)
- [12] D. Kühn and D. Osthus. Matchings in hypergraphs of large minimum degree. *J. Graph Theory*, 51(4):269–280, 2006. [1.2](#)
- [13] D. Kühn, D. Osthus, and A. Treglown. Matchings in 3-uniform hypergraphs. *submitted*. [1.2](#)
- [14] K. Markstrom and A. Ruciński. Perfect matchings and Hamilton cycles in hypergraphs with large degrees. *submitted*, 2010. [1.2](#)
- [15] Ore, Ø. Arc coverings of graphs. *Ann. Mat. Pura Appl., IV. Ser.*, 55:315–321, 1961. [1.1](#)
- [16] O. Pikhurko. Perfect matchings and K_4^3 -tilings in hypergraphs of large codegree. *Graphs Comb.*, 24(4):391–404, 2008. [1.2](#)
- [17] V. Rödl and A. Ruciński. Dirac-type questions for hypergraphs - a survey (or more problems for Endre to solve). *preprint*, 2010. [1.2](#)
- [18] V. Rödl, A. Ruciński, M. Schacht, and E. Szemerédi. A note on perfect matchings in uniform hypergraphs with large minimum collective degree. *Commentat. Math. Univ. Carol.*, 49(4):633–636, 2008. [1.2](#)
- [19] V. Rödl, A. Ruciński, and E. Szemerédi. A Dirac-type theorem for 3-uniform hypergraphs. *Comb. Probab. Comput.*, 15(1-2):229–251, 2006. [1.2](#)
- [20] V. Rödl, A. Ruciński, and E. Szemerédi. Perfect matchings in uniform hypergraphs with large minimum degree. *Eur. J. Comb.*, 27(8):1333–1349, 2006. [1.2](#)
- [21] V. Rödl, A. Ruciński, and E. Szemerédi. An approximate Dirac-type theorem for k -uniform hypergraphs. *Combinatorica*, 28(2):229–260, 2008. [1.2](#)
- [22] V. Rödl, A. Ruciński, and E. Szemerédi. Perfect matchings in large uniform hypergraphs with large minimum collective degree. *J. Comb. Theory, Ser. A*, 116(3):613–636, 2009. [1.2](#)
- [23] V. Rödl, A. Ruciński, and E. Szemerédi. Dirac-type conditions for Hamiltonian paths and cycles in 3-uniform hypergraphs. *submitted*, 2010. [1.2](#)
- [24] Z. Tuza. Steiner system and large non-Hamiltonian hypergraphs. *Matematiche*, 61(1):179–183, 2006. [1.1](#)
- [25] D. Woitas. Permutationsgruppen als Codes - das Bailey-Verfahren. Diploma Thesis, Freie Universität Berlin, 2010. [1.1](#)

INSTITUT FÜR MATHEMATIK, FREIE UNIVERSITÄT BERLIN, ARNIMALLEE 3-5, D-14195 BERLIN, GERMANY
E-mail address: `glebov | weps@mi.fu-berlin.de`

INSTITUT FÜR INFORMATIK, HUMBOLDT-UNIVERSITÄT ZU BERLIN, UNTER DEN LINDEN 6, D-10099 BERLIN, GERMANY
E-mail address: `person@informatik.hu-berlin.de`