

Lifting Galois sections along torsors

NIELS BORNE, MICHEL EMSALEM, AND JAKOB STIX

Abstract — Associated to a smooth curve X/k and an effective étale divisor $D \subseteq X$, we construct torus torsors F_D and E_D over X . If the base field k is \mathbb{Q} and D is a union of torsion packets, we show unconditionally that every Galois section lifts to $\pi_1(F_D)$.

1. INTRODUCTION

Let \bar{k} be a fixed separable closure of an arbitrary field k , and let $\text{Gal}_k = \text{Gal}(\bar{k}/k)$ be the absolute Galois group of k . For a variety X/k , let $X_{\bar{k}} = X \times_k \bar{k}$ be the base change of X to \bar{k} .

1.1. Section conjecture and cuspidalization conjecture. The étale fundamental group $\pi_1(X, \bar{x})$ of X with base point \bar{x} fits into the following exact sequence, denoted by $\pi_1(X/k)$:

$$1 \rightarrow \pi_1(X_{\bar{k}}, \bar{x}) \rightarrow \pi_1(X, \bar{x}) \rightarrow \text{Gal}_k \rightarrow 1. \quad (1.1)$$

By functoriality, any rational point $x \in X(k)$ induces a section $s_X(x) : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$, well defined in the set $\mathcal{S}_{\pi_1(X/k)}$ of sections of (1.1) up to conjugation by an element of $\pi_1(X_{\bar{k}}, \bar{x})$.

In his 1983 anabelian letter to Faltings (see [Gro97]), Grothendieck formulated the *section conjecture*, namely: if k is a field of finite type over \mathbb{Q} , and X is a smooth proper curve of genus at least 2 over k , then the section map

$$s_X : X(k) \rightarrow \mathcal{S}_{\pi_1(X/k)}$$

is a bijective correspondence. He also explained how to deduce the fact that s_X is injective from the Mordell-Weil theorem.

Grothendieck's original vision also contains a version of the section conjecture for a nonempty open subset U of X . It is however easy to see that the direct analogue fails: the section map $s_U : U(k) \rightarrow \mathcal{S}_{\pi_1(U/k)}$ is still injective, but not surjective in general. Indeed for each rational point $x \in (X \setminus U)(k)$ at infinity, one can consider the local extension $\pi_1(U_x/k)$ associated to a punctured formal neighbourhood $U_x = \text{Spec}(\hat{\mathcal{O}}_{X,x}) \setminus \{x\}$ of x in U . This gives rise to a nonempty packet $\mathcal{S}_{\pi_1(U_x/k)} \subset \mathcal{S}_{\pi_1(U/k)}$ of *cuspidal sections*. The generalized section conjecture now asserts that if U is a hyperbolic smooth curve over a field k of finite type over \mathbb{Q} , then sections in $\mathcal{S}_{\pi_1(U/k)}$ either come from rational points of U , or are cuspidal.

The direct observation that the local extensions $\pi_1(U_x/k)$ split has the following striking consequence: if the section conjecture holds for X , then the map $\mathcal{S}_{\pi_1(U/k)} \rightarrow \mathcal{S}_{\pi_1(X/k)}$ is surjective. This consequence of the section conjecture we are fond of calling the *cuspidalization conjecture*.

Besides being a test for the section conjecture, the cuspidalization conjecture is also a significant part of various strategies to prove the section conjecture itself. For example, by working with a fixed and very specific open subset U , namely the complement

$$U = X \setminus X(k)$$

of the set of rational points of X , the cuspidalization conjecture enables to reduce the section conjecture to the statement: if a smooth curve has no rational points, every section is cuspidal. This version of the section conjecture should be more tractable, considering the fact that cuspidal sections have been characterized among all sections by Nakamura (see [Nak90, Sti13]) as those

Date: March 23, 2015.

Niels Borne and Michel Emsalem were supported in part by the Labex CEMPI (ANR-11-LABX-0007-01) and Anr ARIVAF (ANR-10-JCJC 0107).

sections that cyclotomically normalize an inertia subgroup. We emphasize that $X(k) \subseteq X$ is a union of torsion packets in the sense to be described below.

1.2. Main result. Our main result concerns lifting of sections in $\mathcal{S}_{\pi_1(X/k)}$ to the *cuspidally central fundamental group* $\pi_1^{cc}(U)$, which is an intermediate quotient

$$\pi_1(U) \twoheadrightarrow \pi_1^{cc}(U) \twoheadrightarrow \pi_1(X)$$

introduced by Mochizuki (see [Moc07], we recall the original profinite definition in §3.2. Saïdi considered $\pi_1^{cc}(U)$ in the context of the cuspidalization conjecture, see [Saï12]), and we will compare below how Saïdi's approach relates to ours.

The main novelty of this article is that one can relate the geometry of the complement $D = X \setminus U$ to this lifting problem. Following Baker-Poonen (see [BP01]), we will say that a reduced divisor D is a *torsion packet* if geometrically the difference of any two points in the support of D is torsion in the Picard group (see Definition 3). We can now state:

Theorem (Theorem 27). *Let X/\mathbb{Q} be a smooth projective curve of positive genus, let $D \subset X$ be a union of torsion packets, and set $U = X \setminus D$. Then every Galois section $s : \text{Gal}_{\mathbb{Q}} \rightarrow \pi_1(X)$ lifts to a section $\text{Gal}_{\mathbb{Q}} \rightarrow \pi_1^{cc}(U)$.*

We want to stress that Theorem 27 seems to be the first **unconditional** result of lifting to $\pi_1^{cc}(U)$: we make no assumption on the section, in contrast to the setup in [Saï12], which defines "good" sections and shows that these are precisely the ones that can be lifted. An unconditional but weaker result had been obtained by the first two authors in [BE13].

Let us now explain the main ideas of the proof of our result. The first step consists of introducing a torsor F_D over X under the torus $T_D = R_{D/k} \mathbb{G}_m$ whose étale fundamental group $\pi_1(F_D)$ identifies with $\pi_1^{cc}(U)$ (see Proposition 12).

The torsor F_D is itself obtained as a \mathbb{G}_m -torsor over an intermediate torsor E_D over X under the torus $S_D = T_D/\mathbb{G}_m$. In §2, we prove that D is a torsion packet if and only if E_D is torsion (see Proposition 5), a fact that will be crucial in the last step of the proof.

We then study the obstruction of lifting a Galois section along a general torsor E over X under a torus T . We show in Proposition 8 that, if X has $\pi_2^{\text{ét}}(X, \bar{x}) = 0$, the morphism $E \rightarrow X$ gives rise to a natural fibration short exact sequence, denoted by $\pi_1(E/X)$. The next step identifies the class of the extension $\pi_1(E/X)$ with the arithmetic first Chern class $c_1(E)$ of E (see Proposition 16).

Returning to the specific situation of Theorem 27, and given a section s of X , the last and most delicate step consists of killing the obstruction $s^*(c_1(F_D))$. This is the aim of §5. Besides the fact that the torsor E_D is torsion, we use crucially that the relative Brauer group $\text{Br}(X/k) = \ker(\text{Br}(k) \rightarrow \text{Br}(X))$ vanishes when $k = \mathbb{Q}$ in presence of a section, a statement proven by the third author in [Sti10]. It is this result that limits the scope of Theorem 27 to the base field $k = \mathbb{Q}$, a limitation that may be seen as an indication that the section conjecture could be more accessible for the base field \mathbb{Q} (like many other results in arithmetic).

1.3. Notation and conventions. The notation $R_{S'/S}(-)$ denotes Weil restriction of scalars along the finite flat map $S' \rightarrow S$ which is implicit in the notation.

By $X_{S'}$ we will denote the base change $X \times_S S'$ of an S scheme X by $S' \rightarrow S$. However, we would like to direct the kind reader's attention to the following exceptions. The notation S_D, T_D (resp. E_D, F_D) introduced in Section §2 denote a certain torus (resp. torus torsor) associated to a divisor D .

When talking about torsors, one has to fix a topology in which the torsors trivialize locally. This will be the étale topology always without further mention.

For a map $X \rightarrow S$, denote by $\text{Pic}_{X/S}$ the relative Picard scheme. The isomorphism class of a line bundle \mathcal{L} will be denoted by $[\mathcal{L}]$.

Acknowledgments. The authors would like to thank Amaury Thuillier and Angelo Vistoli for discussions and the ENS Lyon for hospitality during a visit of the third author.

2. TORSORS ASSOCIATED TO A DIVISOR ON A CURVE

Let X be a smooth curve defined over $S = \text{Spec}(k)$.

2.1. Definition of torsors E_D and F_D . Let $\Delta : X \rightarrow X \times_S X$ be the diagonal embedding. This is a Cartier divisor, defining an invertible sheaf $\mathcal{O}_{X \times_S X}(\Delta)$. Let D be an effective, étale Cartier divisor on X . We will always assume that $D \neq 0$.

Definition 1. We define torsors

- (1) $F_D = \mathbf{R}_{D \times_S X/X}(\mathbf{Isom}_{D \times_S X}(\mathcal{O}_{X \times_S X}(\Delta)|_{D \times_S X}, \mathcal{O}_{D \times_S X})) \rightarrow X$ under the torus $T_D = \mathbf{R}_{D/S}(\mathbb{G}_m)$,
- (2) $E_D = F_D/\mathbb{G}_m \rightarrow X$ under the quotient torus $S_D = T_D/\mathbb{G}_m$, the cokernel of the adjoint map $\mathbb{G}_m \rightarrow \mathbf{R}_{D/S}\mathbb{G}_m$.

Remark 2. The torsor E_D can also be introduced in a natural way via Picard schemes. Namely, let X_D be the curve obtained from X by pinching D into a single rational point. One can check that the torsor $E_D \rightarrow X$ is the pullback of $\text{Pic}_{X_D/k} \rightarrow \text{Pic}_{X/k}$ via the morphism $X \rightarrow \text{Pic}_{X/k}$ given by Δ . The construction of F_D is somewhat more subtle: one remarks that the morphism $X \rightarrow \text{Pic}_{X/k}$ (thus resp. $E_D \rightarrow \text{Pic}_{X_D/k}$) factors through the Picard stack $\mathcal{P}ic_{X/k}$ (resp. $\mathcal{P}ic_{X_D/k}$). Since the canonical rational point of X_D gives rise to a morphism $\mathcal{P}ic_{X_D/k} \rightarrow \mathbf{B}\mathbb{G}_m$, one gets by composition a morphism $E_D \rightarrow \mathbf{B}\mathbb{G}_m$ and one verifies easily that the corresponding \mathbb{G}_m -torsor is $F_D \rightarrow E_D$.

2.2. Torsion packets and torsion criterion. Let X be a smooth, proper, geometrically connected curve over $S = \text{Spec}(k)$, where k is a field of characteristic 0.

Definition 3 ([BP01]). A (reduced) effective divisor D is a *torsion packet* if any degree 0 divisor on $X_{\bar{k}}$ with support in $D_{\bar{k}}$ is torsion.

Remark 4. (1) Any rational point defines a torsion packet.

- (2) If D is a torsion packet, then any degree 0 divisor on X with support in D is torsion, because $\text{Pic } X \hookrightarrow \text{Pic } X_{\bar{k}}$ is injective.
- (3) However, it is not true that if any degree 0 divisor on X with support in D is torsion, then D is a torsion packet. For instance, if D is irreducible, then any degree 0 divisor on X with support in D is even trivial. But D does not need to be a torsion packet. Indeed, according to [BP01], Corollary 3, if $\text{char } k = 0$ and X is of genus at least 2, the size of torsion packets is bounded. It is thus enough to choose D of degree strictly larger than this size, which is possible if k has an infinite absolute Galois group, to get a counterexample.

Proposition 5. *Let X be a smooth, proper, geometrically connected curve over $S = \text{Spec}(k)$. The torsor $E_D \rightarrow X$ is torsion if and only if the étale divisor $D \subseteq X$ is a torsion packet.*

Proof. The Hochschild-Serre spectral sequence gives the following exact sequence:

$$0 \rightarrow H^1(k, S_D) \rightarrow H^1(X, S_D) \rightarrow H^1(X_{\bar{k}}, S_D)^{\text{Gal}_k}$$

Moreover, the first group is torsion by Lemma 7 below, thus E_D is torsion if and only if $(E_D)_{\bar{k}}$ is torsion. As the formation of E_D also commutes with base change, we can assume that k is algebraically closed. The group of characters $\text{Hom}(S_D, \mathbb{G}_m)$ of S_D is the group $(\bigoplus_{D(k)} \mathbb{Z})^{\Sigma=0}$ of divisors δ of degree 0 with support in D . Since it is free of finite rank, it is enough to show that for such divisor δ , the \mathbb{G}_m -torsor obtained by reduction of the structure group of E_D is torsion. But this \mathbb{G}_m -torsor is nothing else than $\mathbf{Isom}(\mathcal{O}_X, \mathcal{O}_X(\delta))$, and the result follows. \square

Definition 6. (1) For any k -scheme Y , we define the relative Brauer group $\mathrm{Br}(Y/k)$ as the kernel of the natural morphism $\mathrm{Br}(k) \rightarrow \mathrm{Br}(Y)$.

(2) For any k -scheme Y of finite type, we define the index

$$\mathrm{ind}(Y) = \mathrm{gcd} \{ \deg(y) ; y \text{ is a closed point of } Y \},$$

where $\deg(y) = [\kappa(y) : k]$ is the degree of the residue field extension at y .

Lemma 7. *We have*

$$\mathrm{H}^1(k, S_D) = \mathrm{Br}(D/k),$$

and, in particular, $\mathrm{H}^1(k, S_D)$ is torsion and killed by $\mathrm{ind}(D)$.

Proof. This follows from the Galois cohomology exact sequence associated to

$$0 \rightarrow \mathbb{G}_m \rightarrow T_D = \mathrm{R}_{D/S}(\mathbb{G}_m) \rightarrow S_D \rightarrow 0, \quad (2.1)$$

Shapiro's Lemma and Hilbert's Theorem 90. For an extension k'/k and $x \in D(k')$, the pullback

$$x^* : (\mathrm{R}_{D/S}(\mathbb{G}_m))_{k'} \rightarrow \mathbb{G}_{m,k'}$$

splits (2.1) and so

$$\mathrm{Br}(D/k) \hookrightarrow \mathrm{Br}(k'/k)$$

which by a corestriction argument is killed by $[k' : k]$. Hence $\mathrm{Br}(D/k)$ is killed by $\mathrm{ind}(D)$. \square

3. EXTENSIONS OF FUNDAMENTAL GROUPS ASSOCIATED TO TORSORS

3.1. Extensions of fundamental groups associated to fibrations. Following [AM69] and [Fri82] a noetherian scheme X with a geometric point \bar{x} has étale homotopy groups

$$\pi_i^{\mathrm{ét}}(X, \bar{x})$$

for $i \geq 0$. If X is geometrically unibranch, then $\pi_1^{\mathrm{ét}}(X, \bar{x}) = \pi_1(X, \bar{x})$. It is well known that smooth geometrically connected curves X over any field k such that $X_{\bar{k}} \not\cong \mathbb{P}_{\bar{k}}^1$ are algebraic $\mathrm{K}(\pi, 1)$ spaces, which in particular means that $\pi_2^{\mathrm{ét}}(X, \bar{x}) = 0$.

Proposition 8. *Let k be a field of characteristic 0. Let X/k be a geometrically connected and geometrically unibranch variety, let T/k be a torus and $E \rightarrow X$ a torsor under T . Let \bar{y} be a geometric point of E with image $\bar{x} \in X$.*

If $\pi_2^{\mathrm{ét}}(X, \bar{x}) = 0$, then the sequence

$$1 \rightarrow \pi_1(E_{\bar{x}}, \bar{y}) \rightarrow \pi_1(E, \bar{y}) \rightarrow \pi_1(X, \bar{x}) \rightarrow 1 \quad (3.1)$$

is exact.

Proof. Because of the exact sequence (1.1) for E and X , we may assume that k is algebraically closed. Then $T \simeq \mathbb{G}_m^r$ is a trivial torus, and $E \rightarrow X$ is Zariski locally isomorphic to the trivial \mathbb{G}_m^r -torsor. Therefore $E \rightarrow X$ is a geometric fibration in the sense of [Fri82] Definition 11.4. By [Fri82] Theorem 11.5 we have an exact homotopy sequence

$$\pi_2^{\mathrm{ét}}(X, \bar{x}) \rightarrow \pi_1^{\mathrm{ét}}(E_{\bar{x}}, \bar{y}) \rightarrow \pi_1^{\mathrm{ét}}(E, \bar{y}) \rightarrow \pi_1^{\mathrm{ét}}(X, \bar{x}) \rightarrow \pi_0^{\mathrm{ét}}(E_{\bar{x}}, \bar{y})$$

and the claim follows from $\pi_2^{\mathrm{ét}}(X, \bar{x}) = 0$ and $\pi_0^{\mathrm{ét}}(E_{\bar{x}}, \bar{y}) = 0$. \square

Remark 9. The vanishing assumption for $\pi_2^{\mathrm{ét}}(X, \bar{x})$ is indeed important. As an example, we consider an algebraically closed field \bar{k} , some $n \geq 1$, and the \mathbb{G}_m -torsor

$$\mathbb{A}_{\bar{k}}^{n+1} \setminus \{0\} \rightarrow \mathbb{P}_{\bar{k}}^n$$

associated to the line bundle $\mathcal{O}(1)$. Then by Zariski-Nagata purity of the branch locus we have $\pi_1(\mathbb{A}_{\bar{k}}^{n+1} \setminus \{0\}) = 0$ and the sequence

$$\pi_1(\mathbb{G}_{m, \bar{k}}) \rightarrow \pi_1(\mathbb{A}_{\bar{k}}^{n+1} \setminus \{0\}) \rightarrow \pi_1(\mathbb{P}_{\bar{k}}^n) \rightarrow 1$$

is not injective on the left. Indeed, the group $\pi_2^{\mathrm{ét}}(\mathbb{P}_{\bar{k}}^n) \neq 0$ does not vanish.

Remark 10. In the situation of Proposition 8, the choice of \bar{y} defines an isomorphism $T_{\bar{x}} \simeq E_{\bar{x}}$ by translation, and the group

$$\pi_1(T_{\bar{x}}, 1) \simeq \pi_1(E_{\bar{x}}, \bar{y})$$

is the fundamental group of an algebraic group (in characteristic 0, see [Sti13] §13.1) and hence abelian (so that we can neglect base points). The conjugation action of $\pi_1(E, \bar{y})$ on $\pi_1(E_{\bar{x}}, \bar{y})$ thus defines a $\pi_1(X, \bar{x})$ -module structure on $\pi_1(E_{\bar{x}}, \bar{y})$.

Proposition 11. *Let X be a geometrically connected and geometrically unibranch variety over $S = \text{Spec}(k)$ of characteristic 0, and assume that $\pi_2^{\text{ét}}(X, \bar{x}) = 0$. Let T/S be a torus and $E \rightarrow X$ a torsor under T . Then the $\pi_1(X, \bar{x})$ -action on $\pi_1(E_{\bar{x}}, \bar{y})$ factors over the projection $\pi_1(X, \bar{x}) \rightarrow \text{Gal}_k$ and translation is an isomorphism*

$$\tau : \mathbb{T}(T) \simeq \pi_1(T_{\bar{k}}, 1) = \pi_1(T_{X, \bar{x}}, 1) \simeq \pi_1(E_{\bar{x}}, \bar{y})$$

where $\mathbb{T}(T)$ denotes the Tate module of T .

Proof. The $\pi_1(X, \bar{x})$ -module structure on $\pi_1(T_{\bar{k}}, 1) = \pi_1(T_{X, \bar{x}}, 1)$ associated to $T_X \rightarrow X$ comes by functoriality from the action associated to $T \rightarrow \text{Spec}(k)$. The action thus factors through the projection $\pi_1(X, \bar{x}) \rightarrow \text{Gal}_k$. The identification $\mathbb{T}(T) \simeq \pi_1(T_{\bar{k}}, 1)$ as Gal_k -modules is classical, see for example [Sti13] Section §13.1.

We consider the isomorphism $\Phi : T_X \times_X E \rightarrow E \times_X E$ over X defined by multiplication $T \times_X E \rightarrow E$ and second projection $T \times_X E \rightarrow E$. By Proposition 8, we obtain an isomorphism of extensions

$$\begin{array}{ccccccc} 0 & \longrightarrow & \pi_1((T_X \times_X E)_{\bar{x}}, (1, \bar{y})) & \longrightarrow & \pi_1(T_X \times_X E, (1, \bar{y})) & \longrightarrow & \pi_1(X, \bar{x}) \longrightarrow 0 \\ & & \downarrow \simeq & & \downarrow \pi_1(\Phi) & & \downarrow \text{id} \\ 0 & \longrightarrow & \pi_1((E \times_X E)_{\bar{x}}, (\bar{y}, \bar{y})) & \longrightarrow & \pi_1(E \times_X E, (\bar{y}, \bar{y})) & \longrightarrow & \pi_1(X, \bar{x}) \longrightarrow 0 \end{array}$$

Since Φ is a map over E via second projection, we obtain an isomorphism of $\pi_1(X, \bar{x})$ -modules

$$\begin{aligned} \pi_1(T_{\bar{k}}, 1) &= \ker \left(\pi_1((T_X \times_X E)_{\bar{x}}, (1, \bar{y})) \xrightarrow{\pi_1(\text{pr}_2)} \pi_1(E_{\bar{x}}, \bar{y}) \right) \\ &\simeq \ker \left(\pi_1((E \times_X E)_{\bar{x}}, (\bar{y}, \bar{y})) \xrightarrow{\pi_1(\text{pr}_2)} \pi_1(E_{\bar{x}}, \bar{y}) \right) = \pi_1(E_{\bar{x}}, \bar{y}) \end{aligned}$$

induced by Φ . This completes the proof. \square

3.2. Comparison with the maximal cuspidally central quotient. The aim of this paragraph is to mention the following interpretation of $\pi_1(F_D, \bar{y})$. Let $U = X \setminus D$ be the complement of the support of the divisor and set

$$N = \text{Ker}(\pi_1(U, \bar{x}) \rightarrow \pi_1(X, \bar{x})).$$

Recall the notion of the maximal **cuspidally central** quotient $\pi_1^{\text{cc}}(U, \bar{x})$ due to Mochizuki [Moc07] Definition 1.5(i): the biggest quotient $\pi_1^{\text{cc}}(U, \bar{x}) = \pi_1(U, \bar{x})/N_{\text{cc}}$ by a normal subgroup $N_{\text{cc}} \subseteq N$ such that one gets an exact sequence

$$1 \rightarrow N^{\text{cc}} \rightarrow \pi_1^{\text{cc}}(U, \bar{x}) \rightarrow \pi_1(X, \bar{x}) \rightarrow 1$$

where $N^{\text{cc}} = N/N_{\text{cc}}$ is abelian, and the action of $\pi_1(X_{\bar{k}}, \bar{x})$ by conjugation on N^{cc} is trivial.

Proposition 12. *The canonical lift $U \rightarrow F_D$ of the inclusion $U \subset X$ induces an isomorphism*

$$\pi_1^{\text{cc}}(U, \bar{x}) \simeq \pi_1(F_D, \bar{x}).$$

Proof. If D is totally split, this follows from [Moc07], Proposition 1.8 iii). Since the claim is of geometric nature, the general case follows. \square

4. OBSTRUCTIONS TO LIFTING GALOIS SECTIONS ALONG TORSORS UNDER TORI

In this section, we fix a smooth, proper, geometrically connected curve X of genus at least 1 over a field k of characteristic 0, and T/k a torus. Let $E \rightarrow X$ be a torsor under the torus T . Then E is also geometrically connected and therefore defines a fundamental exact sequence analogous to (1.1). The issue we want to address is: does a given section $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ of (1.1) lift to a section $\text{Gal}_k \rightarrow \pi_1(E, \bar{y})$?

4.1. Arithmetic first Chern class. We introduce in this paragraph the notion of arithmetic first Chern class of a torsor under a torus. The relevance of this notion for anabelian issues has been pointed out by Mochizuki in the case of line bundles (see [Moc99] Definition 0.3). Our definition is the straightforward generalization.

The most logical way to proceed is to use Jannsen's cohomology (see [Jan88]), that is, cohomology defined on the category of inverse systems of étale sheaves of abelian groups on the small étale site of the base scheme X . In this context, the Kummer short exact sequence takes the following form, for a pair of integers (m, n) such that $m|n$:

$$\begin{array}{ccccccc} 0 & \longrightarrow & T[n] & \longrightarrow & T & \xrightarrow{n \cdot} & T \longrightarrow 0 \\ & & \downarrow \frac{n \cdot}{m \cdot} & & \downarrow \frac{n \cdot}{m \cdot} & & \downarrow \text{id} \\ 0 & \longrightarrow & T[m] & \longrightarrow & T & \xrightarrow{m \cdot} & T \longrightarrow 0 \end{array} \quad (4.1)$$

Let us define $\mathbb{T}(T)$ (resp. $(T, \frac{n \cdot}{m \cdot})$) as the inverse system given by the left (resp. middle) column of diagram (4.1). Note that the Jannsen cohomology of the right column is the usual étale cohomology of T .

Definition 13. The *first arithmetic Chern class* of a T -torsor $E \rightarrow X$ is the image of its class by the coboundary morphism

$$c_1 : H^1(X, T) \rightarrow H^2(X, \mathbb{T}(T))$$

in the Jannsen cohomology long exact sequence associated to the short exact sequence (4.1).

Proposition 14.

(1) *The following morphism is an isomorphism:*

$$H^2(X, \mathbb{T}(T)) \rightarrow \varprojlim_{n \in \mathbb{N}} H^2(X, T[n]).$$

(2) *For a torus T over a field k , the following sequence is exact:*

$$0 \rightarrow H^1(k, T) \xrightarrow{c_1} H^2(k, \mathbb{T}(T)) \rightarrow \mathbb{T}(H^2(k, T)) \rightarrow 0.$$

Proof. (1) The short exact \varprojlim -sequence (see [Jan88], (3.1))

$$0 \rightarrow \varprojlim_{n \in \mathbb{N}}^1 H^1(X, T[n]) \rightarrow H^2(X, \mathbb{T}(T)) \rightarrow \varprojlim_{n \in \mathbb{N}} H^2(X, T[n]) \rightarrow 0$$

reduces assertion (1) to the vanishing of the \varprojlim^1 -term. The Kummer sequence provides the exact sequence

$$0 \rightarrow T(X)/nT(X) \rightarrow H^1(X, T[n]) \rightarrow H^1(X, T)[n] \rightarrow 0,$$

which by Mittag-Leffler and $\varprojlim_{n \in \mathbb{N}}^2 = 0$ leads to an exact sequence

$$0 = \varprojlim_{n \in \mathbb{N}}^1 T(X)/nT(X) \rightarrow \varprojlim_{n \in \mathbb{N}}^1 H^1(X, T[n]) \xrightarrow{\sim} \varprojlim_{n \in \mathbb{N}}^1 H^1(X, T)[n] \rightarrow 0.$$

The restriction to n -torsion of the short exact sequence of low degree terms of the Hochschild–Serre spectral sequence for $X \rightarrow \text{Spec}(k)$ and coefficients T yields exactness of

$$0 \rightarrow H^1(k, T)[n] \rightarrow H^1(X, T)[n] \rightarrow H^1(X_{\bar{k}}, T)[n]^{\text{Gal}_k}.$$

By Hilbert's Theorem 90, the groups $H^1(k, T)[n]$ have bounded exponent and thus the system $\mathbb{T}(H^1(k, T))$ is Mittag-Leffler zero ([Jan88], 1.10). Since $T_{\bar{k}} \cong \mathbb{G}_m^d$ for $d = \dim T$ we find non-canonically

$$H^1(X_{\bar{k}}, T) \cong \text{Pic}(X_{\bar{k}})^d$$

which has finite n -torsion. Thus the projective system $(H^1(X, T)[n])$ is an extension of a system of finite levels

$$\left(\text{im}(H^1(X, T)[n] \rightarrow H^1(X_{\bar{k}}, T)[n]^{\text{Gal}_k}) \right)$$

by a Mittag-Leffler zero system. Therefore its $\varprojlim_{n \in \mathbb{N}}^1$ vanishes and the proof is complete.

(2) The Kummer sequences for T induces exact sequences

$$0 \rightarrow H^1(k, T)/n H^1(k, T) \rightarrow H^2(k, T[n]) \rightarrow H^2(k, T)[n] \rightarrow 0.$$

Since by Hilbert's Theorem 90 the group $H^1(k, T)$ has bounded exponent, we have

$$\begin{aligned} \varprojlim_n H^1(k, T)/n H^1(k, T) &= H^1(k, T) \\ \varprojlim_n^1 H^1(k, T)/n H^1(k, T) &= 0. \end{aligned}$$

Therefore assertion (2) follows by passing to the limit and (1) for $X = \text{Spec}(k)$. \square

Remark 15. Because the morphism $H^2(X, \mathbb{T}(T)) \rightarrow \varprojlim_{n \in \mathbb{N}} H^2(X, T[n])$ is an isomorphism, it will be often sufficient to consider only Chern classes modulo n , denoted by $c_1(E)_n$, and defined as the image of $c_1(E)$ by the morphism $H^2(X, \mathbb{T}(T)) \rightarrow H^2(X, T[n])$.

4.2. Class of the fibration. Let $h : E \rightarrow X$ be a T -torsor, and consider the exact sequence

$$1 \rightarrow \pi_1(T_{\bar{k}}, 1) \rightarrow \pi_1(E, \bar{y}) \rightarrow \pi_1(X, \bar{x}) \rightarrow 1 \quad (4.2)$$

from Proposition 8 where we use the isomorphism $\pi_1(T_{\bar{k}}, 1) \simeq \pi_1(E_{\bar{x}}, \bar{y})$ of Proposition 11. This yields an abelian cohomology class

$$\pi_1(E/X) \in H^2(\pi_1(X, \bar{x}), \pi_1(T_{\bar{k}}, 1)).$$

The section s lifts to (a Galois section of) E if and only if $s^*(\pi_1(E/X)) = 0$ in $H^2(\text{Gal}_k, \pi_1(T_{\bar{k}}, 1))$. We are therefore interested in an explicit description of the class $\pi_1(E/X)$.

Proposition 16. *Let X be a geometrically connected and geometrically unibranch variety over a field k of characteristic 0. Assume that $\pi_2^{\text{ét}}(X, \bar{x}) = 0$.*

Then the morphism $H^2(\pi_1(X, \bar{x}), \pi_1(T_{\bar{k}}, 1)) \rightarrow H^2(X, \mathbb{T}(T))$ is an isomorphism and sends $\pi_1(E/X)$ to $c_1(E)$.

Proof. Let $\tilde{X}_{\text{ét}}$ be the universal covering of the étale homotopy type $X_{\text{ét}}$ with respect to the geometric point $\bar{x} \in X$. Then for any locally constant constructible torsion sheaf \mathcal{F} we have isomorphisms

$$\begin{aligned} H^0(\tilde{X}_{\text{ét}}, \mathcal{F}) &\simeq \mathcal{F}_{\bar{x}}, \\ H^1(\tilde{X}_{\text{ét}}, \mathcal{F}) &= 0, \\ H^2(\tilde{X}_{\text{ét}}, \mathcal{F}) &\simeq \text{Hom}(\pi_2^{\text{ét}}(X, \bar{x}), \mathcal{F}_{\bar{x}}). \end{aligned}$$

The exact sequence of low degree terms for the Leray spectral sequence for $\tilde{X}_{\text{ét}} \rightarrow X_{\text{ét}}$ yields

$$H^1(\pi_1(X, \bar{x}), \mathcal{F}_{\bar{x}}) \simeq H^1(X, \mathcal{F})$$

and the exact sequence

$$0 \rightarrow H^2(\pi_1(X, \bar{x}), \mathcal{F}_{\bar{x}}) \rightarrow H^2(X, \mathcal{F}) \rightarrow \text{Hom}_{\pi_1(X, \bar{x})}(\pi_2^{\text{ét}}(X, \bar{x}), \mathcal{F}_{\bar{x}}).$$

For $\mathcal{F} = T[n]$, using the short exact \varprojlim -sequence (see [Jan88], (3.1)) shows that

$$H^2(\pi_1(X, \bar{x}), \pi_1(T_{\bar{k}}, 1)) \simeq H^2(\pi_1(X, \bar{x}), \mathbb{T}(T)) \rightarrow H^2(X, \mathbb{T}(T))$$

is an isomorphism.

We now prove the second claim based on ideas by Mochizuki [Moc03] Lemma 4.4+5. Denote by $\text{Et}(X)$ the small étale site of X , and by $\text{FEt}(X)$ the finite étale site of X . The natural morphism $\gamma : \text{Et}(X) \rightarrow \text{FEt}(X)$ induces for each sheaf \mathcal{F} on $\text{FEt}(X)$ and the corresponding locally constant sheaf $\gamma^*\mathcal{F}$ on $\text{Et}(X)$ and each $i \geq 0$ a morphism

$$\mathrm{H}^i(\pi_1(X, \bar{x}), \mathcal{F}_{\bar{x}}) \rightarrow \mathrm{H}^i(X, \gamma^*\mathcal{F})$$

via the usual identification of sheaves on $\text{FEt}(X)$ and representations of $\pi_1(X, \bar{x})$. Note that this coincides with the edge map of the spectral sequence used above.

Now the fibration $E \rightarrow X$ gives rise to a commutative diagram of sites:

$$\begin{array}{ccc} \text{Et}(E) & \longrightarrow & \text{FEt}(E) \\ \downarrow & & \downarrow \\ \text{Et}(X) & \longrightarrow & \text{FEt}(X) \end{array}$$

This diagram induces in turn a morphism of Leray-Serre spectral sequences associated to the sheaf $\mathbb{T}(T)$ corresponding to the $\pi_1(E)$ -representation $\pi_1(T_{\bar{k}}, 1)$ (we omit base points for readability sake):

$$\begin{array}{ccc} \mathrm{H}^p(\pi_1(X), \mathrm{H}^q(\pi_1(T_{\bar{k}}), \pi_1(T_{\bar{k}}))) & \longrightarrow & \mathrm{H}^p(X, \mathrm{H}^q(T_{\bar{k}}, \mathbb{T}(T))) \\ \Downarrow & & \Downarrow \\ \mathrm{H}^{p+q}(\pi_1(E), \pi_1(T_{\bar{k}})) & \longrightarrow & \mathrm{H}^{p+q}(E, \mathbb{T}(T)) \end{array}$$

The transgression morphisms $d_2^{0,1} : E_2^{0,1} \rightarrow E_2^{2,0}$ are thus compatible in the sense that the following diagram commutes:

$$\begin{array}{ccc} \text{End}_{\text{Gal}_k}(\pi_1(T_{\bar{k}})) & \xrightarrow{\cong} & \text{End}_k(\mathbb{T}(T)) \\ \downarrow d_2^{0,1} & & \downarrow d_2^{0,1} \\ \mathrm{H}^2(\pi_1(X), \pi_1(T_{\bar{k}})) & \longrightarrow & \mathrm{H}^2(X, \mathbb{T}(T)) \end{array}$$

The top arrow is induced by $\pi_1(T_{\bar{k}}) \simeq \mathbb{T}(T)$, in particular it sends id to id . Now the profinite version of [HS53, III, Theorem 4] implies that the left vertical map sends id to $-\pi_1(E/X)$ and a similar argument with Čech cocycles shows that the right vertical map sends id to $-c_1(E)$. \square

4.3. Killing torsion obstructions. By Proposition 16 we associate to a T -torsor $E \rightarrow X$ and a section $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ a class

$$s^*(c_1(E)) := s^*(\pi_1(E/X)) \in \mathrm{H}^2(k, \mathbb{T}(T)).$$

Lemma 17. $s^*(c_1(E)) = 0$ if and only if s lifts to $\pi_1(E, \bar{y})$.

Proof. This follows from Proposition 16. \square

We can make this obstruction more tractable thanks to the following lemma.

Lemma 18. *The Tate module $\mathbb{T}(A)$ of an abelian group A is torsion free.*

Proof. This is clear from the expression $\mathbb{T}(A) = \text{Hom}(\mathbb{Q}/\mathbb{Z}, A)$. \square

If $E \rightarrow X$ is torsion, so is $s^*(c_1(E))$, and Proposition 14 together with Lemma 18 show that the obstruction $s^*(c_1(E))$ lives in fact in $\mathrm{H}^1(k, T)$.

5. LIFTING TO F_D OVER THE RATIONALS

In this section, k will be a field of characteristic 0 and X/k a smooth, projective, geometrically connected curve of genus ≥ 1 . Let $D \subset X$ be an effective reduced Cartier divisor and $U = X \setminus D$. We consider the associated torsors $F_D \rightarrow X$ and $E_D \rightarrow X$ from Definition 1 and study the lifting obstruction for sections $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ to the fundamental group of the respective torsors.

5.1. Vanishing of Brauer obstructions. The short exact sequence of low degree terms of the Leray spectral sequence for $X \rightarrow \text{Spec}(k)$ and \mathbb{G}_m reads

$$0 \rightarrow \text{Pic}(X) \rightarrow \text{Pic}_{X/k}(k) \xrightarrow{b} \text{Br}(k) \rightarrow \text{Br}(X). \quad (5.1)$$

By definition the map b is the Brauer obstruction map with values in the relative Brauer group $\text{Br}(X/k)$, see Definition 6, that measures the failure of a rational point of the Picard variety to describe an actual line bundle.

Proposition 19. *Let X/k be a smooth projective curve of positive genus such that $\pi_1(X/k)$ admits a section. Then the following holds.*

- (1) $b(L) = 0$ for all torsion points $L \in \text{Pic}_{X/k}(k)_{\text{tors}}$.
- (2) If k/\mathbb{Q}_p is a finite extension, then $\#\text{Br}(X/k)$ is a power of p .
- (3) If $k = \mathbb{R}$, then $\text{Br}(X/k) = 0$.

Proof. Assertion (1) is [Sti10] Proposition 12, and (2) is proven in [Sti10] Theorem 15.

Assertion (3) follows from the real section conjecture, see [Sti13] §16.1. Indeed, any section $s : \text{Gal}_{\mathbb{R}} \rightarrow X$ comes from a point $x \in X(\mathbb{R})$, and evaluation in x yields a retraction to $\text{Br}(k) \rightarrow \text{Br}(X)$, showing $\text{Br}(X/k) = 0$. \square

Corollary 20. *Let X/\mathbb{Q} be a smooth projective curve of positive genus such that $\pi_1(X/\mathbb{Q})$ admits a section. Then the relative Brauer group $\text{Br}(X/\mathbb{Q})$ vanishes.*

Proof. The Hasse–Brauer–Noether theorem shows that

$$\text{Br}(X/\mathbb{Q}) \hookrightarrow \ker \left(\bigoplus_v \text{Br}(X \times_{\mathbb{Q}} \mathbb{Q}_v/\mathbb{Q}_v) \xrightarrow{\sum_v \text{inv}_v} \mathbb{Q}/\mathbb{Z} \right)$$

is injective, where v ranges over all places of \mathbb{Q} . Base change of sections implies that for all places v of \mathbb{Q} the extension $\pi_1(X \times_{\mathbb{Q}} \mathbb{Q}_v/\mathbb{Q}_v)$ splits. By Proposition 19 then $\text{Br}(X \times_{\mathbb{Q}} \mathbb{R}/\mathbb{R}) = 0$ and $\text{Br}(X \times_{\mathbb{Q}} \mathbb{Q}_p/\mathbb{Q}_p)$ is cyclic of p -power order. This forces $\text{Br}(X/\mathbb{Q}) = 0$. \square

5.2. Divisibility of line bundles. Let us recall the definition of a neighbourhood of a section.

Definition 21. A **neighbourhood** of a section $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ is a connected finite étale cover $h : X' \rightarrow X$ together with a lift

$$s' : \text{Gal}_k \rightarrow \pi_1(X', \bar{x}') \subseteq \pi_1(X, \bar{x})$$

of s . A short notation for a neighbourhood is (X', s') .

Neighbourhoods are geometrically connected over k , because $\pi_1(X', \bar{x}') \rightarrow \text{Gal}_k$ is surjective.

Example 22. A wealth of neighbourhoods are constructed as follows. Let $\varphi : \pi_1(X_{\bar{k}}, \bar{x}) \rightarrow G$ be a characteristic finite quotient. Then $\ker(\varphi)$ is a normal subgroup in $\pi_1(X, \bar{x})$ and

$$\pi_1(X_{\varphi}, \bar{x}) = \langle \ker(\varphi), s(\text{Gal}_k) \rangle = \{ \gamma s(\sigma) ; \varphi(\gamma) = 1, \sigma \in \text{Gal}_k \} \subseteq \pi_1(X, \bar{x})$$

together with the obvious lift describes a neighbourhood of s . Moreover, we have $\pi_1(X_{\varphi, \bar{k}}, \bar{x}) = \ker(\varphi)$, so that

$$\deg(X_{\varphi} \rightarrow X) = \#G.$$

Since $\pi_1(X_{\bar{k}}, \bar{x})$ is topologically finitely generated, the neighbourhoods X_{φ} form a cofinal system in the system $X_s = (X')$ of all neighbourhoods.

Proposition 23. *Let X/k be a smooth projective curve of positive genus. Let $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ be a section and let \mathcal{L} be a line bundle on X . Then the following holds.*

- (1) *For every $n \geq 1$ there is a neighbourhood (X', s') of s , such that there is a $M \in \text{Pic}_{X'}(k)$ with*

$$[\mathcal{L}|_{X'}] = M^{\otimes n}.$$

- (2) *If, moreover, for every neighbourhood (X', s') of s , the relative Brauer group $\text{Br}(X'/k)$ vanishes, then for every $n \geq 1$ there is a neighbourhood (X', s') of s , such that there is a line bundle \mathcal{M} on X' with*

$$\mathcal{L}|_{X'} \simeq \mathcal{M}^{\otimes n}.$$

Definition 24. If the conclusion (2) of Proposition 23 holds, then we say that the **line bundle \mathcal{L} is divisible locally in neighbourhoods of s** .

Proof of Proposition 23. Let $\text{Pic}_{X/k}^{n*}$ denote the subgroup of the Picard variety of line bundles of degree divisible by n . The boundary map to

$$0 \rightarrow \text{Pic}_{X/k}[n] \rightarrow \text{Pic}_{X/k} \rightarrow \text{Pic}_{X/k}^{n*} \rightarrow 0,$$

namely

$$\text{Pic}_{X/k}^{n*}(k) \rightarrow \text{H}^1(k, \text{Pic}_{X/k}[n]),$$

describes the obstruction to being divisible by n in the Picard variety for line bundles of degree divisible by n . This obstruction is natural under pullback.

We first prove assertion (1). Since $n \mid \#\pi_1(X_{\bar{k}}, \bar{x})$ we find a neighbourhood (X_1, s_1) of s with $n \mid \deg(X_1/X)$. Then $[\mathcal{L}|_{X_1}] \in \text{Pic}_{X_1/k}^{n*}(k)$ because

$$\deg(\mathcal{L}|_{X_1, \bar{k}}) = \deg(X_1/X) \cdot \deg(\mathcal{L}).$$

Let (X_2, s_2) be the neighbourhood of s_1 associated to the maximal abelian quotient of exponent n of $\pi_1(X_{1, \bar{k}}, \bar{x}_1)$. Then the induced map

$$\text{Pic}_{X_1/k}[n] = \text{Hom}(\pi_1(X_{1, \bar{k}}, \bar{x}_1), \mathbb{Z}/n\mathbb{Z}(1)) \rightarrow \text{Hom}(\pi_1(X_{2, \bar{k}}, \bar{x}_2), \mathbb{Z}/n\mathbb{Z}(1)) = \text{Pic}_{X_2/k}[n]$$

is the zero map. Thus the obstruction for $[\mathcal{L}|_{X_1}]$ to divisibility by n in the Picard variety vanishes after restriction to X_2 . This proves (1).

Let $M \in \text{Pic}_{X_2/k}(k)$ be an n th root of $[\mathcal{L}|_{X_2}]$. In order to prove (2) we have to investigate the Brauer obstruction for M to come from an actual line bundle. But this is the class $b(M)$ for the map b in (5.1) for X_2/k and b vanishes by assumption. This concludes the proof of (2). \square

Proposition 25. *Let X/k be a smooth projective curve of positive genus, let $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ be a Galois section, and let \mathcal{L} be a line bundle on X . Then \mathcal{L} is locally divisible in neighbourhoods of s if and only if $s^*(c_1(\mathcal{L})) = 0$.*

Proof. Let X_s be the projective limit of the pro-system of all neighbourhoods of s . Then \mathcal{L} is locally divisible in neighbourhoods of s if and only if $\mathcal{L}|_{X_s}$ is divisible in $\text{Pic}(X_s)$.

The Kummer sequence on X_s yields the exact sequence

$$\text{Pic}(X_s) \xrightarrow{n} \text{Pic}(X_s) \xrightarrow{c_1(\cdot)_n} \text{H}^2(X_s, \mathbb{Z}/n\mathbb{Z}(1)),$$

so that $\mathcal{L}|_{X_s}$ is divisible by n on X_s if and only if $c_1(\mathcal{L}|_{X_s})_n = 0$.

Naturality of the first Chern class and the isomorphism

$$\text{H}^2(X_s, \mathbb{Z}/n\mathbb{Z}(1)) = \text{H}^2(\pi_1(X_s, \bar{x}), \mathbb{Z}/n\mathbb{Z}(1)) \xrightarrow{s^*} \text{H}^2(k, \mathbb{Z}/n\mathbb{Z}(1))$$

show that $c_1(\mathcal{L}|_{X_s})_n = 0$ if and only if $s^*(c_1(\mathcal{L})_n) = 0$. Moreover, $s^*(c_1(\mathcal{L})_n) = 0$ for all $n \geq 1$ if and only if $s^*(c_1(\mathcal{L})) = 0$, because by Proposition 14 (1) for $T = \mathbb{G}_m$ and $X = \text{Spec}(k)$ we have

$$\text{H}^2(k, \hat{\mathbb{Z}}(1)) \simeq \varprojlim_{n \in \mathbb{N}} \text{H}^2(k, \mathbb{Z}/n\mathbb{Z}(1)). \quad \square$$

Proposition 26. *Let X/k be a smooth projective curve of positive genus, and let $s : \text{Gal}_k \rightarrow \pi_1(X, \bar{x})$ be a Galois section. Consider the following assertions.*

- (a) *All line bundles \mathcal{L} on X are locally divisible in neighbourhoods of s .*
- (b) *$s^* \circ c_1 : \text{Pic}(X) \rightarrow \text{H}^2(k, \widehat{\mathbb{Z}}(1))$ vanishes.*
- (c) *The relative Brauer group $\text{Br}(X/k)$ vanishes.*
- (c') *The relative Brauer group $\text{Br}(X'/k)$ vanishes for all neighbourhoods X' of s .*

Then the following implications hold:

$$(c') \implies (a) \iff (b) \implies (c).$$

Proof. (c') \implies (a) was proven in Proposition 23 (2). The equivalence of (a) with (b) follows from Proposition 25.

For (a) \implies (c) we have to show that $b(L) = 0$ for all $L \in \text{Pic}_{X/k}(k)$. Since $b(L) \in \text{Br}(X/k)$ is torsion, there is an $n \geq 1$ such that $L^{\otimes n} = [\mathcal{M}]$ for a line bundle \mathcal{M} on X . By assumption (a), there is a neighbourhood (X', s') of s such that $\mathcal{M}|_{X'}$ admits an n th root

$$\mathcal{M}|_{X'} = \mathcal{L}'^{\otimes n}$$

with a line bundle \mathcal{L}' on X' . The difference $\Delta = L|_{X'} - [\mathcal{L}']$ is an n -torsion element in $\text{Pic}_{X'/k}(k)$. By Proposition 19 (1) we compute

$$b(L) = b(L|_{X'}) = b(\Delta) + b([\mathcal{L}']) = 0,$$

and this proves (c). \square

5.3. Lifting to F_D over \mathbb{Q} . In this section we study the lifting problem over the field \mathbb{Q} .

Theorem 27. *Let X/\mathbb{Q} be a smooth projective curve of positive genus, and let $D \subset X$ be a union of torsion packets. Then every Galois section $s : \text{Gal}_{\mathbb{Q}} \rightarrow \pi_1(X, \bar{x})$ lifts to a section $\text{Gal}_{\mathbb{Q}} \rightarrow \pi_1(F_D, \bar{y})$.*

Proof. Let $D = \bigcup_{i=1}^n D_i$ be the decomposition into torsion packets $D_i \subset X$. The torsor F_D is the product (over X) of the F_{D_i} , and similarly $\pi_1(F_D)$ is the fibre product (over $\pi_1(X)$) of the $\pi_1(F_{D_i})$, so we can assume that D is a single torsion packet.

The section s lifts if and only if $s^*(c_1(F_D)) = 0$. Since T_D is the restriction of scalars of \mathbb{G}_m , Shapiro's Lemma and Hilbert's Theorem 90 imply $\text{H}^1(k, T_D) = 0$. Using Proposition 14 and Lemma 18, we find that $s^*(c_1(F_D))$ takes values in the torsion free group

$$\text{H}^2(k, \mathbb{T}(T_D)) \simeq \mathbb{T}(\text{H}^2(k, T_D)).$$

Hence, we may replace F_D by a multiple. The result follows from a diagram chase in

$$\begin{array}{ccccc}
 & & \text{H}^1(X, \mathbb{G}_m) & \xrightarrow{c_1} & \text{H}^2(X, \widehat{\mathbb{Z}}(1)) \\
 & \swarrow i_* & & \searrow & \downarrow s^* \\
 & & \text{H}^1(X, T_D) & \xrightarrow{c_1} & \text{H}^2(X, \mathbb{T}(T_D)) \\
 & & & \downarrow s^* & \\
 & & & & \text{H}^2(k, \widehat{\mathbb{Z}}(1)) \\
 & & & \swarrow i_* & \\
 0 & \longrightarrow & \text{H}^1(k, T_D) = 0 & \longrightarrow & \text{H}^2(k, \mathbb{T}(T_D)) \longrightarrow \mathbb{T}(\text{H}^2(k, T_D))
 \end{array}$$

Namely, since D is a torsion packet, we can apply Proposition 5 and choose an integer $N \geq 1$ so that $N \cdot E_D$ is trivial. Then $N \cdot F_D$ comes from a \mathbb{G}_m -torsor, and the result follows from Proposition 26 and Corollary 20. \square

Corollary 28. *Let X/\mathbb{Q} be a smooth projective curve of positive genus, and let $U = X \setminus X(\mathbb{Q})$ be the complement of the set of all \mathbb{Q} -rational points. Then every Galois section $s : \text{Gal}_{\mathbb{Q}} \rightarrow \pi_1(X, \bar{x})$ lifts to a section $\text{Gal}_{\mathbb{Q}} \rightarrow \pi_1^{\text{cc}}(U, \bar{x})$.*

Proof. If $X(\mathbb{Q})$ is infinite (at most for X of genus 1), then we understand $\pi_1^{\text{cc}}(U, \bar{x})$ as the natural projective limit of $\pi_1^{\text{cc}}(X \setminus D, \bar{x})$ with D ranging over all finite subsets $D \subseteq X(\mathbb{Q})$. We may therefore restrict to the case of $U = X \setminus D$ and $D \subseteq X(\mathbb{Q})$ a finite set.

The divisor $D \subset X$ is a union of torsion packets. Therefore the section s lifts to a section $\text{Gal}_{\mathbb{Q}} \rightarrow \pi_1(F_D, \bar{y})$ by Theorem 27, and since $\pi_1^{\text{cc}}(U, \bar{x}) \simeq \pi_1(F_D, \bar{y})$ by Proposition 12, this completes the proof. \square

Remark 29. Note that similar arguments enable to show over a local field k a prime to p version of Theorem 27, with $\pi_1(F_D, \bar{y})$ replaced by its quotient by the p -part of the geometric fundamental group $\pi_1((T_D)_{\bar{k}})$.

REFERENCES

- [AM69] Michel Artin and Barry Mazur. *Étale homotopy*, volume 100 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 1969.
- [BE13] Niels Borne and Michel Emsalem. Un critère d'épointage des sections l -adiques. *To appear in Bulletin de la Société Mathématique de France*, 2013.
- [BP01] Matthew Baker and Bjorn Poonen. Torsion packets on curves. *Compositio Mathematica*, 127(1):109–116, 2001.
- [Fri82] Eric M Friedlander. *Étale homotopy of simplicial schemes*, volume 104 of *Annals of Mathematical Studies*. Princeton University Press, 1982.
- [Gro97] Alexander Grothendieck. Brief an G. Faltings. In *Geometric Galois actions, 1*, volume 242 of *London Math. Soc. Lecture Note Ser.*, pages 49–58. Cambridge Univ. Press, Cambridge, 1997. With an English translation on pp. 285–293.
- [HS53] G. Hochschild and J.-P. Serre. Cohomology of group extensions. *Trans. Amer. Math. Soc.*, 74:110–134, 1953.
- [Jan88] Uwe Jannsen. Continuous étale cohomology. *Mathematische Annalen*, 280(2):207–245, 1988.
- [Moc99] Shinichi Mochizuki. The local pro- p anabelian geometry of curves. *Invent. Math.*, 138(2):319–423, 1999.
- [Moc03] Shinichi Mochizuki. Topics surrounding the anabelian geometry of hyperbolic curves. In *Galois groups and fundamental groups*, volume 41 of *Math. Sci. Res. Inst. Publ.*, pages 119–165. Cambridge Univ. Press, Cambridge, 2003.
- [Moc07] Shinichi Mochizuki. Absolute anabelian cuspidalizations of proper hyperbolic curves. *J. Math. Kyoto Univ.*, 47(3):451–539, 2007.
- [Nak90] Hiroaki Nakamura. Galois rigidity of the étale fundamental groups of punctured projective lines. *J. Reine Angew. Math.*, 411:205–216, 1990.
- [Sai12] Mohamed Saïdi. The cuspidalisation of sections of arithmetic fundamental groups. *Adv. Math.*, 230(4-6):1931–1954, 2012.
- [Sti10] Jakob Stix. On the period-index problem in light of the section conjecture. *Amer. J. Math.*, 132(1):157–180, 2010.
- [Sti13] Jakob Stix. *Rational Points and Arithmetic of Fundamental Groups: Evidence for the Section Conjecture*, volume 2054 of *Lecture Notes in Mathematics*. Springer-Verlag, Berlin, 2013.

NIELS BORNE, U.M.R. CNRS 8524, U.F.R. DE MATHÉMATIQUES, UNIVERSITÉ LILLE 1, 59 655 VILLENEUVE D'ASCQ CÉDEX, FRANCE

E-mail address: Niels.Borne@math.univ-lille1.fr

MICHEL EMSALEM, U.M.R. CNRS 8524, U.F.R. DE MATHÉMATIQUES, UNIVERSITÉ LILLE 1, 59 655 VILLENEUVE D'ASCQ CÉDEX, FRANCE

E-mail address: Michel.Emsalem@math.univ-lille1.fr

JAKOB STIX, INSTITUT FÜR MATHEMATIK, JOHANN WOLFGANG GOETHE-UNIVERSITÄT FRANKFURT, ROBERT-MAYER-STRASSE 6–8, 60325 FRANKFURT AM MAIN, GERMANY

E-mail address: stix@math.uni-frankfurt.de